

10/620209

=> d his

(FILE 'HOME' ENTERED AT 17:57:21 ON 16 JAN 2004)

FILE 'CAPLUS' ENTERED AT 18:00:09 ON 16 JAN 2004
E SCHLOEMER G/IN

L1 17 S E4-E6
L2 0 S L1 AND ACETAMIDE?
L3 0 S L1 AND IMIDAZO?
L4 7 S L1 AND PROCESS
L5 5 S DIMETHYLGLYCOXYLAMIDE
SELECT L5 5 RN

L6 FILE 'REGISTRY' ENTERED AT 18:16:26 ON 16 JAN 2004
3 S E1-E3

FILE 'CAPLUS' ENTERED AT 18:18:30 ON 16 JAN 2004
SELECT L5 4 RN

L7 FILE 'REGISTRY' ENTERED AT 18:18:53 ON 16 JAN 2004
3 S E4-E6

L8 FILE 'REGISTRY' ENTERED AT 18:30:24 ON 16 JAN 2004
STRUCTURE UPLOADED
L9 0 S L8

L10 FILE 'BEILSTEIN' ENTERED AT 18:31:02 ON 16 JAN 2004
0 S L8
L11 1 S L8 SSS FULL


L12 FILE 'REGISTRY' ENTERED AT 18:32:54 ON 16 JAN 2004
2 S L8 SSS FULL

L13 FILE 'CAPLUS' ENTERED AT 18:33:36 ON 16 JAN 2004
4 S L12

=> d l8

L8 HAS NO ANSWERS

L8 STR




G1 Me, Et, n-Pr, i-Pr, n-Bu, i-Bu, s-Bu, t-Bu


10/620209

=> d 1-4 bib abs hitstr

L13 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN
AN 1991:101100 CAPLUS
DN 114:101100
TI One-pot synthesis of N,N,N',N'-tetrasubstituted ureas and oxomalonamides by oxidative carbonylation of lithium amides at atmospheric pressure
AU Nudelman, Norma S.; Lewkowicz, Elizabeth S.; Perez, Daniel G.
CS Fac. Cienc. Exactas, Univ. Buenos Aires, Buenos Aires, 1428, Argent.
SO Synthesis (1990), (10), 917-20
CODEN: SYNTBF; ISSN: 0039-7881
DT Journal
LA English
OS CASREACT 114:101100
AB N,N,N',N'-tetrasubstituted ureas RR1NCONRR1 (R = R1 = Bu, cyclohexyl, CHMe2, cyclohexyl) were prepd. in good yields by reaction of lithium aliph. amides RR1NLi in THF soln. with CO under mild conditions (0.degree., 1013 mbar) followed by treatment with oxygen prior to work up. N,N,N',N'-tetrasubstituted oxomalonamides (oxopropanediamides) RR1NCOCONRR1 were prepd. under similar reaction conditions by carrying out the reaction in the presence of known amts. of the pure amine. Besides being an useful synthetic method, the present studies afford new evidence of the mechanism of the reaction.
IT 83862-73-1P
RL: SPN (Synthetic preparation); PREP (Preparation)
(prepn. of)
RN 83862-73-1 CAPLUS
CN Acetamide, 2,2'-oxybis[N,N-dibutyl-2-hydroxy- (9CI) (CA INDEX NAME)




L13 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN
AN 1988:55481 CAPLUS
DN 108:55481
TI Carbon-carbon bond formation through the carbonylation of lithium dialkylamides. One-pot synthesis of N-alkyl-substituted formamides, glyoxylamides, and hydroxymalonamides
AU Perez, Daniel G.; Nudelman, N. Sbarbati
CS Fac. Cienc. Exactas, Univ. Buenos Aires, Buenos Aires, 1428, Argent.
SO Journal of Organic Chemistry (1988), 53(2), 408-13
CODEN: JOCEAH; ISSN: 0022-3263
DT Journal
LA English
OS CASREACT 108:55481
AB The reaction of RR1NLi (R = R1 = Bu, pentyl, cyclohexyl; R = iso-Pr, R1 = cyclohexyl; RR1 = 3-oxapentamethylene) with CO to yield RR1NCHO, (RR1NCOCHOH)2O, and RR1NCOCH(OH)CONRR1 (R, R1 = same as above) was examd. under a no. of different conditions. Evidence supporting a lithium carbamoyl intermediate for the latter compds. is presented. A general procedure for the prepn. of tetraalkylureas, tetraalkyloxalamides, and tetraalkyloxomalonamides is given.
IT 83862-73-1P
RL: SPN (Synthetic preparation); PREP (Preparation)
(prepn. of)
RN 83862-73-1 CAPLUS
CN Acetamide, 2,2'-oxybis[N,N-dibutyl-2-hydroxy- (9CI) (CA INDEX NAME)




L13 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN
AN 1983:34210 CAPLUS
DN 98:34210
TI Insertion of carbon monoxide into lithium-nitrogen bonds. One-pot synthesis of dialkylformamides and dialkylglyoxylamides
AU Nudelman, N. Sbarbati; Perez, Daniel
CS Fac. Cienc. Exactas Nat., Univ. Buenos Aires, Buenos Aires, 1428, Argent.
SO Journal of Organic Chemistry (1983), 48(1), 133-4
CODEN: JOCEAH; ISSN: 0022-3263

10/620209


DT Journal
LA English
OS CASREACT 98:34210
AB Lithium dialkylamides react with CO to afford dialkylformamides (I),
tetraalkylhydroxymalonamides and dialkylglyoxylamides (II). Reaction
conditions are described to produce I or II in good yields.
IT 83862-73-1P
RL: SPN (Synthetic preparation); PREP (Preparation)
(prepn. of)
RN 83862-73-1 CAPLUS
CN Acetamide, 2,2'-oxybis[N,N-dibutyl-2-hydroxy- (9CI) (CA INDEX NAME)



L13 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN
AN 1979:121688 CAPLUS
DN 90:121688
TI Carbonylation of lithium dialkylamides to carbamoyllithium derivatives
AU Rautenstrauch, Valentin; Joyeux, Michel
CS Forschungslab., Firmenich S. A., Geneva, Switz.
SO Angewandte Chemie (1979), 91(1), 72-3
CODEN: ANCEAD; ISSN: 0044-8249
DT Journal
LA German
OS CASREACT 90:121688
GI



AB Lithiation of piperidine gave lithium piperidide, carbonylation of which
gave I which was hydrolyzed to give 20-30% II. Reaction of I with
cyclohexanone gave 85% 4:1 III-IV. Similarly, (Me2CH)2NH was lithiated,
carbonylated, and hydrolyzed to give 20-40% (Me2CH)2NCHO.
IT 68986-67-4P
RL: SPN (Synthetic preparation); PREP (Preparation)
(prepn. of)
RN 68986-67-4 CAPLUS
CN Acetamide, 2,2'-oxybis[2-hydroxy-N,N-bis(1-methylethyl)- (9CI) (CA INDEX
NAME)



10/620209

=> s dimethylglyoxylamide
L5 5 DIMETHYLGLYYOXYLAMIDE

=> d 1-5 bib abs kwic

L5 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN
AN 2003:887680 CAPLUS
DN 139:364844
TI Preparation of indolizines as sPLA2 inhibitors
IN Dillard, Robert D.; Hagishita, Sanji; Ohtani, Mitsuaki
PA Eli Lilly and Company, USA; Shiongi and Company, Ltd.
SO U.S., 62 pp., Cont.-in-part of U.S. Ser. No. 278,445.
CODEN: USXXAM
DT Patent
LA English
FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6645976	B1	20031111	US 1997-765566	19970428
	WO 9603383	A1	19960208	WO 1995-US9381	19950720
	W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT				
	RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
PRAI	US 1994-278445	A2	19940721		
	WO 1995-US9381	W	19950720		
OS	MARPAT 139:364844				
GI					

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. I, II, III [wherein X = O or S; R11 = independently H, alkyl, or halo; R12 = H, halo, (cyclo)alkyl, cycloalkenyl, alkoxy, alkylthio, or a non-interfering substituent having 1-3 atoms other than H; R13 = (un)substituted alkyl, alkenyl, alkynyl, (hetero)cyclyl optionally connected by a linking group; R15 and R16 = independently H, non-interfering substituent, or (un)substituted (hetero)cyclyl; R17 and R18 = independently H, non-interfering substituent, or acidic linker; with the proviso that at least one of R17 and R18 must be an acidic linker; or pharmaceutically acceptable salt, ester, or amide prodrug derivs. thereof], and their 3-acetamide, 3-acetic acid hydrazide, and 3-glyoxylamide analogs were prep'd. as inhibitors of human secreted phospholipase A2 (sPLA2) mediated release of fatty acids. For example, conversion of 2-methyl-5-methoxypyridine to the anion in THF using lithium diisopropylamide and subsequent reaction with benzonitrile produced 5-methoxy-2-phenacylpyridine (57.0%). Cyclization of the pyridine deriv. with 1-bromo-2-butanone using NaHCO3 in acetone gave the 1-benzoylindolizine (90.7%), which was reduced by LAH to give 1-benzyl-2-ethyl-6-methoxyindolizine (94.5%). Acylation (98.5%) with Et oxalyl chloride in benzene, followed by sapon. with LiOH in H2O and amidation using NH4OH, provided 2-(1-benzyl-2-ethyl-6-methoxyindolizin-3-yl)glyoxylamide. Demethylation by BBr3 in CH2Cl2, coupling with Et 4-bromobutyrate (56.2%) in the presence of NaH in DMF, and hydrolysis with LiOH gave the title indolizine IV (49.9%). Eighty-eight compds. of the invention inhibited recombinant human sPLA2 in a chromogenic assay with IC50 values ranging from 0.006 .mu.M to 1.1 .mu.M, in contrast to IC50 values >50 .mu.M for comparative examples. Administration of 10/mg/kg of the representative compd., 2-[8-(carbomethoxymethoxy)-2-ethyl-3-(2-phenylbenzyl)indolizin-1-yl]glyoxylamide, improved the survival rate of male Wistar rats with sPLA2-induced pancreatitis from 33.3% (vehicle) to 91.7%. Thus, invention compds. and their pharmaceutical formulations are useful for the treatment of conditions such as septic shock, adult respiratory distress syndrome, pancreatitis, trauma, bronchial asthma, allergic rhinitis, and rheumatoid arthritis.

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

IT 177556-77-3P, 2-(3-Benzyl-8-hydroxy-2-ethylindolizin-1-yl)acetamide
177556-79-5P, 2-[2-Ethyl-8-hydroxy-3-(o-phenylbenzyl)indolizin-1-yl]acetamide 177556-80-8P, 2-[3-(m-Chlorobenzyl)-2-ethyl-8-hydroxyindolizin-1-yl]acetamide 177556-81-9P, 2-[2-Cyclopropyl-8-hydroxy-3-(o-phenylbenzyl)indolizin-1-yl]acetamide 177556-84-2P,
2-[8-[[(Benzylloxycarbonyl)methyl]oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-

1-yl)acetamide 177556-85-3P, 2-[8-[[[(Benzyloxycarbonyl)methyl]oxy]-3-(m-chlorobenzyl)-2-ethylindolizin-1-yl]acetamide 177556-86-4P,
 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-cyclopropyl-3-(o-phenylbenzyl)indolizin-1-yl]acetamide 177556-92-2P, 2-(3-Benzyl-2-methylindolizin-1-yl)glyoxylamide 177556-93-3P, 2-(3-Benzyl-8-benzyloxy-2-ethylindolizin-1-yl)glyoxylamide 177556-94-4P, 2-(3-Benzyl-8-benzyloxy-2-ethylindolizin-1-yl)-N-methylglyoxylamide 177556-95-5P, 2-(3-Benzyl-8-benzyloxy-2-ethylindolizin-1-yl)-N,N-dimethylglyoxylamide 177556-96-6P,
 2-[8-Benzyloxy-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 177556-97-7P, 2-[8-Benzyloxy-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]-N-methylglyoxylamide 177556-98-8P, 2-[8-Benzyloxy-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]-N,N-dimethylglyoxylamide 177556-99-9P, 2-(3-Benzyl-8-benzyloxy-2-methylindolizin-1-yl)glyoxylamide 177557-00-5P, 2-[8-Benzyloxy-3-(m-chlorobenzyl)-2-ethylindolizin-1-yl]glyoxylamide 177557-01-6P, 2-[8-Benzyloxy-2-ethyl-3-(m-trifluoromethylbenzyl)indolizin-1-yl]glyoxylamide 177557-02-7P,
 2-[8-Benzyloxy-2-ethyl-3-(1-naphthylmethyl)indolizin-1-yl]glyoxylamide 177557-03-8P, 2-[8-Benzyloxy-2-cyclopropyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 177557-25-4P, 2-[8-Benzyloxy-2-ethyl-3-(p-phenylbenzyl)indolizin-1-yl]glyoxylamide 177557-26-5P,
 2-(8-Benzyloxy-3-cyclohexylmethyl-2-ethylindolizin-1-yl)glyoxylamide 177557-27-6P, 2-(8-Benzyloxy-3-cyclopentylmethyl-2-ethylindolizin-1-yl)glyoxylamide 177557-28-7P, 2-(8-Benzyloxy-3-cycloheptylmethyl-2-ethylindolizin-1-yl)glyoxylamide 177557-29-8P, 2-[8-Benzyloxy-2-ethyl-3-pentylindolizin-1-yl]glyoxylamide 177557-30-1P, 2-[8-Benzyloxy-2-ethyl-3-(2-propylpentyl)indolizin-1-yl]glyoxylamide 177557-31-2P,
 2-[8-Benzyloxy-2-ethyl-3-[(naphth-2-yl)methyl]indolizin-1-yl]glyoxylamide 177557-32-3P, 2-[8-Benzyloxy-3-(3,5-di-tert-butylbenzyl)-2-ethylindolizin-1-yl]glyoxylamide 177557-33-4P, 2-[8-Benzyloxy-2-ethyl-3-(2-phenylethyl)indolizin-1-yl]glyoxylamide 177557-34-5P,
 2-[8-Benzyloxy-3-(o-benzylbenzyl)-2-ethylindolizin-1-yl]glyoxylamide 177557-35-6P, 2-[8-Benzyloxy-2-ethyl-3-[(thiophen-2-yl)methyl]indolizin-1-yl]glyoxylamide 177557-36-7P, 2-[8-Benzyloxy-2-ethyl-3-[[3-(thiophen-2-yl)thiophen-2-yl]methyl]indolizin-1-yl]glyoxylamide 177557-37-8P,
 2-[8-Benzyloxy-2-ethyl-3-(m-methoxybenzyl)indolizin-1-yl]glyoxylamide 177557-38-9P, 2-[8-Benzyloxy-2-ethyl-3-(o-nitrobenzyl)indolizin-1-yl]glyoxylamide 177557-41-4P, 2-[8-Benzyloxy-3-[(biphenyl-2-yl)methyl]-2-methylindolizin-1-yl]glyoxylamide 177557-42-5P, 2-(3-Benzyl-8-benzyloxy-2-cyclopropylindolizin-1-yl)glyoxylamide 177557-43-6P,
 2-[3-(p-Butylbenzyl)-2-ethyl-8-methoxyindolizin-1-yl]glyoxylamide 177557-44-7P, 2-(8-Benzyloxy-3-cyclohexylmethyl-2-methylindolizin-1-yl)glyoxylamide 177557-48-1P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-3-cyclohexylmethyl-2-ethylindolizin-1-yl]glyoxylamide 177557-49-2P,
 2-[8-[[[(Carbomethoxy)methyl]oxy]-3-cyclopentylmethyl-2-ethylindolizin-1-yl]glyoxylamide 177557-50-5P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-3-cycloheptylmethyl-2-ethylindolizin-1-yl]glyoxylamide 177557-51-6P,
 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-pentylindolizin-1-yl]glyoxylamide 177557-52-7P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-(2-propylpentyl)indolizin-1-yl]glyoxylamide 177557-53-8P,
 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-[(naphth-2-yl)methyl]indolizin-1-yl]glyoxylamide 177557-55-0P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-(2-phenylethyl)indolizin-1-yl]glyoxylamide 177557-56-1P,
 2-[3-(o-Benzylbenzyl)-8-[[[(Carbomethoxy)methyl]oxy]-2-ethylindolizin-1-yl]glyoxylamide 177557-58-3P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-[[3-(thiophen-2-yl)thiophen-2-yl]methyl]indolizin-1-yl]glyoxylamide 177557-59-4P, 2-[2-Ethyl-8-[[[(Carbomethoxy)methyl]oxy]-3-(m-methoxybenzyl)indolizin-1-yl]glyoxylamide 177557-60-7P,
 2-[2-Ethyl-8-[[[(Carbomethoxy)methyl]oxy]-3-(o-nitrobenzyl)indolizin-1-yl]glyoxylamide 177557-62-9P, 2-[3-[(Adamant-1-yl)methyl]-8-[[[(Carbomethoxy)methyl]oxy]-2-methylindolizin-1-yl]glyoxylamide 177557-63-0P, 2-[3-Benzyl-8-[[[(Carbomethoxy)methyl]oxy]-2-cyclopropylindolizin-1-yl]glyoxylamide 177557-64-1P,
 2-[3-(p-Butylbenzyl)-8-[[[(Carbomethoxy)methyl]oxy]-2-ethylindolizin-1-yl]glyoxylamide 177557-65-2P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-3-cyclohexylmethyl-2-methylindolizin-1-yl]glyoxylamide 177557-74-3P,
 2-[8-[[[(Carboxymethyl)oxy]-3-(3,5-di-tert-butylbenzyl)-2-ethylindolizin-1-yl]glyoxylamide 177557-88-9P, 2-[3-[(Biphenyl-2-yl)methyl]-8-[[[(Carbomethoxy)methyl]oxy]-2-methylindolizin-1-yl]glyoxylamide 177557-89-0P, 2-[3-[(Biphenyl-2-yl)methyl]-8-[[[(Carbomethoxy)methyl]oxy]-2-isopropylindolizin-1-yl]glyoxylamide 177557-90-3P, 2-[2-tert-Butyl-8-[[[(Carbomethoxy)methyl]oxy]-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 177557-91-4P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-cyclopentyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 177557-92-5P,
 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-(m-phenylbenzyl)indolizin-1-yl]glyoxylamide 177557-93-6P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-(3-phenyl-2-propenyl)indolizin-1-yl]glyoxylamide 177557-94-7P,
 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-(3-phenylpropyl)indolizin-1-yl]glyoxylamide 177557-95-8P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-

cyclopropyl-3-[(1-naphthyl)methyl]indolizin-1-yl]glyoxylamide
 177557-96-9P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-3-cyclohexylmethyl-2-
 cyclopropylindolizin-1-yl]glyoxylamide 177557-97-0P,
 2-[3-[(Biphenyl-2-yl)methyl]-8-(carboxymethoxy)-2-methylindolizin-1-
 yl]glyoxylamide 177557-99-2P, 2-[2-tert-Butyl-8-(carboxymethoxy)-3-(o-
 phenylbenzyl)indolizin-1-yl]glyoxylamide 177558-00-8P,
 2-[8-[(Carboxymethyl)oxy]-2-cyclopentyl-3-(o-phenylbenzyl)indolizin-1-
 yl]glyoxylamide 177558-01-9P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(m-
 phenylbenzyl)indolizin-1-yl]glyoxylamide 177558-02-0P,
 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(3-phenyl-2-propenyl)indolizin-1-
 yl]glyoxylamide 177558-03-1P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(3-
 phenylpropyl)indolizin-1-yl]glyoxylamide 177558-04-2P,
 2-[8-(Carboxymethoxy)-2-cyclopropyl-3-[(1-naphthyl)methyl]indolizin-1-
 yl]glyoxylamide 177558-05-3P, 2-[8-(Carboxymethoxy)-3-cyclohexylmethyl-2-
 cyclopropylindolizin-1-yl]glyoxylamide 177558-34-8P,
 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-methylthio-3-(o-phenylbenzyl)indolizin-
 1-yl]glyoxylamide 177558-90-6P, 2-(3-Benzyl-2-ethyl-8-hydroxyindolizin-1-
 yl]glyoxylamide 177558-91-7P, 2-[2-Ethyl-8-hydroxy-3-(o-
 phenylbenzyl)indolizin-1-yl]glyoxylamide 177558-92-8P,
 2-[2-Ethyl-8-hydroxy-3-(o-phenylbenzyl)indolizin-1-yl]-N-
 methylglyoxylamide 177558-93-9P, 2-[2-Ethyl-8-hydroxy-3-(o-
 phenylbenzyl)indolizin-1-yl]-N,N-dimethylglyoxylamide
 177558-94-0P, 2-(3-Benzyl-8-hydroxy-2-methylindolizin-1-yl]glyoxylamide
 177558-95-1P, 2-[3-(m-Chlorobenzyl)-2-ethyl-8-hydroxyindolizin-1-
 yl]glyoxylamide 177558-96-2P, 2-[2-Ethyl-8-hydroxy-3-(m-
 trifluoromethylbenzyl)indolizin-1-yl]glyoxylamide 177558-97-3P,
 2-[2-Ethyl-8-hydroxy-3-[(1-naphthyl)methyl]indolizin-1-yl]glyoxylamide
 177558-98-4P, 2-[2-Cyclopropyl-8-hydroxy-3-(o-phenylbenzyl)indolizin-1-
 yl]glyoxylamide 177559-41-0P, 2-[2-Ethyl-8-hydroxy-3-[(thiophen-2-
 yl)methyl]indolizin-1-yl]glyoxylamide 177559-42-1P, 2-[2-Ethyl-8-hydroxy-
 3-(m-methoxybenzyl)indolizin-1-yl]glyoxylamide 177559-43-2P,
 2-[2-Ethyl-8-hydroxy-3-(o-nitrobenzyl)indolizin-1-yl]glyoxylamide
 177559-45-4P, 2-[3-[(Biphenyl-2-yl)methyl]-8-hydroxy-2-methylindolizin-1-
 yl]glyoxylamide 177559-46-5P, 2-(3-Benzyl-2-cyclopropyl-8-
 hydroxyindolizin-1-yl]glyoxylamide 177559-47-6P, 2-[3-(p-Butylbenzyl)-2-
 ethyl-8-hydroxyindolizin-1-yl]glyoxylamide 177559-79-4P,
 2-[3-Benzyl-7-(3-carbethoxypropyloxy)-2-ethylindolizin-1-yl]glyoxylamide
 177559-80-7P, 2-[7-(3-Carbethoxypropyloxy)-2-cyclopropyl-3-(o-
 phenylbenzyl)indolizin-1-yl]glyoxylamide 177559-81-8P,
 2-[7-(3-Carbethoxypropyloxy)-2-ethyl-3-(o-phenylbenzyl)indolizin-1-
 yl]glyoxylamide 177559-82-9P, 2-[7-[[3-(Carbethoxy)propyl]oxy]-3-
 cyclohexylmethyl-2-ethylindolizin-1-yl]glyoxylamide 177559-84-1P,
 2-[7-[[[(Carboethoxy)methyl]oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-1-
 yl]glyoxylamide 177559-99-8P, 2-[3-Benzyl-8-
 [[(methoxycarbonyl)methyl]amino]-2-methylindolizin-1-yl]glyoxylamide
 177560-00-8P, 2-[8-[[[(Methoxycarbonyl)methyl]amino]-3-cyclohexylmethyl-2-
 methylindolizin-1-yl]glyoxylamide 182115-63-5P, Methyl
 2-[[3-Benzyl-1-(carbamoylmethyl)-2-ethylindolizin-8-yl]oxy]acetate
 182115-76-0P, 2-[8-Benzoyloxy-2-ethyl-3-[(4-pentylcyclohexyl)methyl]indoliz-
 in-1-yl]glyoxylamide 182115-78-2P, 2-[8-Benzoyloxy-3-cyclopentylmethyl-2-
 cyclopropylindolizin-1-yl]glyoxylamide 182115-84-0P,
 2-[8-Hydroxy-2-ethyl-3-(4-pentylcyclohexylmethyl)indolizin-1-
 yl]glyoxylamide 182115-86-2P, 2-[3-Benzyl-8-[[[(carbethoxy)methyl]oxy]-2-
 ethylindolizin-1-yl]glyoxylamide 182115-87-3P, 2-[8-
 [[[(Carbethoxy)methyl]oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-1-
 yl]glyoxylamide 182115-88-4P, 2-[8-[[[(Carbethoxy)methyl]oxy]-3-(m-
 chlorobenzyl)-2-ethylindolizin-1-yl]glyoxylamide 182115-90-8P,
 2-[8-[[[(Carbethoxy)methyl]oxy]-2-ethyl-3-[(1-naphthyl)methyl]indolizin-1-
 yl]glyoxylamide 182115-92-0P, 2-[3-Benzyl-8-[[[(carbethoxy)methyl]oxy]-2-
 methylindolizin-1-yl]glyoxylamide 182115-93-1P, 2-[8-
 [[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-[(4-pentylcyclohexyl)methyl]indolizi-
 n-1-yl]glyoxylamide 182116-42-3P, 2-[7-(5-Carboethoxypentyloxy)-2-ethyl-
 3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 182116-44-5P,
 2-[3-Benzyl-8-[[[(methoxycarbonyl)methyl]amino]-2-methylindolizin-1-
 yl]acetamide 182116-45-6P, 2-[3-Benzyl-8-[(carboxymethyl)amino]-2-
 methylindolizin-1-yl]acetamide 182116-49-0P, 2-[8-(3-
 Carbomethoxypropyloxy)-2-ethyl-3-(o-phenylbenzyl)indolizin-1-
 yl]glyoxylamide 215160-62-6P, 2-[8-[[[(Carbomethoxy)methyl]oxy]-2-ethyl-3-
 (o-phenylbenzyl)indolizin-1-yl]glyoxylamide 215160-63-7P,
 2-[3-Benzyl-8-[[[(tert-butoxycarbonyl)methyl]oxy]-2-ethylindolizin-1-
 yl]glyoxylamide 215160-64-8P 215160-65-9P 622835-99-8P,
 2-[3-(1-Naphthyl)-8-hydroxy-2-ethylindolizin-1-yl]acetamide
 622836-00-4P, Methyl 2-[[3-Naphthyl-1-(carbamoylmethyl)-2-ethylindolizin-8-
 yl]oxy]acetate 622836-03-7P, 2-[3-Benzyl-8-[[[(carbethoxy)methyl]oxy]-2-
 ethylindolizin-1-yl]-N-methylglyoxylamide 622836-04-8P,
 2-[3-Benzyl-8-[[[(carbethoxy)methyl]oxy]-2-ethylindolizin-1-yl]-N,N-
 dimethylglyoxylamide 622836-05-9P, 2-[8-
 [[[(Carbethoxy)methyl]oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]-N,N-

dimethylglyoxylamide 622836-07-1P, 2-[8-(Cyanomethyloxy)-2-ethyl-3-[(1-naphthyl)methyl]indolizin-1-yl]glyoxylamide 622836-33-3P, 2-[3-[(Adamant-1-yl)methyl]-8-benzoyloxy-2-ethylindolizin-1-yl]glyoxylamide 622836-34-4P, 8-Benzoyloxy-3-(cyclopentylcarbonyl)-2-cyclopropylindolizine 622836-35-5P, 8-Benzoyloxy-3-cyclopentylmethyl-2-cyclopropylindolizine 622836-36-6P, 2-[8-[[[Carbomethoxy)methyl]oxy]-2-ethyl-3-(thiophen-2-yl)indolizin-1-yl]glyoxylamide 622836-37-7P, 2-(3-Cyclopentylmethyl-2-cyclopropyl-8-hydroxyindolizin-1-yl)glyoxylamide 622836-57-1P, 2-[3-(Biphenyl 2-yl)-8-[[[carbomethoxy)methyl]oxy]-2-methoxyindolizin-1-yl]glyoxylamide
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (sPLA2 inhibitor; prepn. of indolizines as inhibitors of sPLA2 mediated release of fatty acids)

IT 177556-76-2P, 2-[1-Benzyl-6-(3-carboxypropyloxy)-2-ethylindolizin-3-yl]glyoxylamide 177556-87-5P, 2-[3-Benzyl-8-[(carboxymethyl)oxy]-2-ethylindolizin-1-yl]acetamide 177556-89-7P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]acetamide 177556-90-0P, 2-[8-[(Carboxymethyl)oxy]-3-(m-chlorobenzyl)-2-ethylindolizin-1-yl]acetamide 177556-91-1P, 2-[8-[(Carboxymethyl)oxy]-2-cyclopropyl-3-(o-phenylbenzyl)indolizin-1-yl]acetamide 177557-67-4P, 2-[2-Ethyl-8-[(carboxymethyl)oxy]-3-(p-phenylbenzyl)indolizin-1-yl]glyoxylamide 177557-68-5P, 2-[8-[(Carboxymethyl)oxy]-3-cyclohexylmethyl-2-ethylindolizin-1-yl]glyoxylamide 177557-69-6P, 2-[8-[(Carboxymethyl)oxy]-3-cyclopentylmethyl-2-ethylindolizin-1-yl]glyoxylamide 177557-70-9P, 2-[8-[(Carboxymethyl)oxy]-3-cycloheptylmethyl-2-ethylindolizin-1-yl]glyoxylamide 177557-71-0P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-pentylindolizin-1-yl]glyoxylamide 177557-72-1P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(2-propylpentyl)indolizin-1-yl]glyoxylamide 177557-75-4P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(2-phenylethyl)indolizin-1-yl]glyoxylamide 177557-76-5P, 2-[8-[(Carboxymethyl)oxy]-3-(o-benzylbenzyl)-2-ethylindolizin-1-yl]glyoxylamide 177557-77-6P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-[(thiophen-2-yl)methyl]indolizin-1-yl]glyoxylamide 177557-78-7P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-[[3-(thiophen-2-yl)thiophen-2-yl]methyl]indolizin-1-yl]glyoxylamide 177557-79-8P, 2-[2-Ethyl-8-[(carboxymethyl)oxy]-3-(m-methoxybenzyl)indolizin-1-yl]glyoxylamide 177557-80-1P, 2-[2-Ethyl-8-[(carboxymethyl)oxy]-3-(o-nitrobenzyl)indolizin-1-yl]glyoxylamide 177557-82-3P, 2-[3-[(Adamant-1-yl)methyl]-8-[(carboxymethyl)oxy]-2-methylindolizin-1-yl]glyoxylamide 177557-83-4P, 2-[3-Benzyl-8-[(carboxymethyl)oxy]-2-cyclopropylindolizin-1-yl]glyoxylamide 177557-84-5P, 2-[3-(p-Butylbenzyl)-8-[(carboxymethyl)oxy]-2-ethylindolizin-1-yl]glyoxylamide 177557-85-6P, 2-[8-[(Carboxymethyl)oxy]-3-cyclohexylmethyl-2-methylindolizin-1-yl]glyoxylamide 177557-86-7P, 2-[8-[(Carboxymethyl)oxy]-3-cyclopentylmethyl-2-cyclopropylindolizin-1-yl]glyoxylamide 177557-87-8P, 2-[8-[(Carboxymethyl)oxy]-3-cyclopentylmethyl-2-methylindolizin-1-yl]glyoxylamide 177558-06-4P, 2-[3-[(Biphenyl-2-yl)methyl]-8-[(carboxymethyl)oxy]-2-ethylindolizin-1-yl]glyoxylamide sodium salt 177558-07-5P, 2-[3-[(Biphenyl-2-yl)methyl]-8-[[[1-(methoxycarbonyloxy)ethoxy]carbonyl]methoxy]-2-ethylindolizin-1-yl]glyoxylamide 177558-08-6P, 2-[3-[(Biphenyl-2-yl)methyl]-2-ethyl-8-[[[1-(isopropylloxycarbonyloxy)ethoxy]carbonyl]methoxy]indolizin-1-yl]glyoxylamide 177558-11-1P, 2-[3-[(Biphenyl-2-yl)methyl]-8-[[[[[1-(cyclopentylcarbonyl)oxy]ethyl]oxy]carbonyl]methyl]oxy]-2-ethylindolizin-1-yl]glyoxylamide 177558-12-2P, 2-[3-[(Biphenyl-2-yl)methyl]-8-[[[[[1-(cyclopentylcarbonyl)oxy]ethyl]oxy]carbonyl]methyl]oxy]-2-ethylindolizin-1-yl]glyoxylamide 177558-18-8P, 2-[3-[(Biphenyl-2-yl)methyl]-2-ethyl-8-[[[1H-tetrazol-5-yl]methyl]oxy]indolizin-1-yl]glyoxylamide 177558-22-4P, 2-[3-Benzyl-7-(3-carboxypropyloxy)-2-ethylindolizin-1-yl]glyoxylamide 177558-23-5P, 2-[7-(3-Carboxypropyloxy)-2-cyclopropyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 177558-24-6P, 2-[7-(3-Carboxypropyloxy)-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 177558-25-7P, 2-[7-(3-Carboxypropyloxy)-3-cyclohexylmethyl-2-ethylindolizin-1-yl]glyoxylamide 177558-26-8P, 2-[3-[(Biphenyl-2-yl)methyl]-8-(3-carboxypropyloxy)-2-ethylindolizin-1-yl]glyoxylamide 177558-27-9P, 2-[7-[(Carboxymethyl)oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 177558-29-1P, 2-[3-[(Biphenyl-2-yl)methyl]-8-(2-carboxyethyloxy)-2-ethylindolizin-1-yl]glyoxylamide 177558-31-5P, 2-[3-[(Biphenyl-2-yl)methyl]-8-(2-carbomethoxyethyloxy)-2-ethylindolizin-1-yl]glyoxylamide 177558-32-6P, 2-[7-(3-Carboxypropyloxy)-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]acetamide 177558-33-7P, 2-[7-(3-Carboxypropyloxy)-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]acetamide 177558-35-9P, 2-[8-[(Carboxymethyl)oxy]-2-methylthio-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 177560-01-9P, 2-[3-Benzyl-8-[(carboxymethyl)amino]-2-methylindolizin-1-yl]glyoxylamide 177560-02-0P, 2-[8-

[(Carboxymethyl)amino]-3-cyclohexylmethyl-2-methylindolizin-1-ylglyoxylamide 182115-96-4P, 2-[3-Benzyl-8-[(carboxymethyl)oxy]-2-ethylindolizin-1-yl]glyoxylamide 182115-97-5P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 182115-98-6P, 2-[8-[(Carboxymethyl)oxy]-3-(m-chlorobenzyl)-2-ethylindolizin-1-yl]glyoxylamide 182115-99-7P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(m-trifluoromethylbenzyl)indolizin-1-yl]glyoxylamide 182116-00-3P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-[(1-naphthyl)methyl]indolizin-1-yl]glyoxylamide 182116-01-4P, 2-[8-[(Carboxymethyl)oxy]-2-cyclopropyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 182116-02-5P, 2-[3-Benzyl-8-[(carboxymethyl)oxy]-2-methylindolizin-1-yl]glyoxylamide 182116-03-6P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(4-pentylcyclohexylmethyl)indolizin-1-yl]glyoxylamide 182116-43-4P, 2-[7-(5-Carboxypentyl)oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 182116-46-7P, 2-[3-[(Biphenyl-2-yl)methyl]-2-methyl-8-[(pyridin-2-yl)methoxy]indolizin-1-yl]glyoxylamide 182116-47-8P, 2-[3-[(Biphenyl-2-yl)methyl]-2-methyl-8-[(pyridin-4-yl)methoxy]indolizin-1-yl]glyoxylamide 182116-48-9P, 2-[3-[(Biphenyl-2-yl)methyl]-2-methyl-8-[(quinolin-2-yl)methoxy]indolizin-1-yl]glyoxylamide 182116-50-3P, 2-[3-Benzyl-8-[(carboxymethyl)oxy]-2-ethylindolizin-1-yl]-N-methylglyoxylamide 182116-51-4P, 2-[3-Benzyl-8-[(carboxymethyl)oxy]-2-ethylindolizin-1-yl]-N,N-dimethylglyoxylamide 622836-01-5P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(1-naphthyl)indolizin-1-yl]acetamide 622836-06-0P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]-N,N-dimethylglyoxylamide 622836-08-2P, 2-[8-[(1H-Tetrazol-5-yl)methyl]oxy]-2-ethyl-3-[(1-naphthyl)methyl]indolizin-1-yl]glyoxylamide 622836-38-8P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(naphth-2-yl)indolizin-1-yl]glyoxylamide 622836-39-9P, 2-[8-[(Carboxymethyl)oxy]-2-ethyl-3-(4-pentylcyclohexylmethyl)indolizin-1-yl]glyoxylamide sodium salt 622836-40-2P, 2-[8-[(Carboxymethyl)oxy]-3-cyclohexylmethyl-2-methylindolizin-1-yl]glyoxylamide sodium salt 622836-41-3P, 2-[8-[(Carboxymethyl)oxy]-3-cyclopentylmethyl-2-methylindolizin-1-yl]glyoxylamide sodium salt 622836-43-5P, 2-[3-[(Biphenyl-2-yl)methyl]-8-[[[[(tert-butoxycarbonyl)methyl]oxy]carbonyl]methyl]oxy]-2-ethylindolizin-1-yl]glyoxylamide 622836-44-6P, 2-[3-[(Biphenyl-2-yl)methyl]-8-[[[[(1-cyclohexyloxy)carbonyl]ethyl]oxy]carbonyl]methyl]oxy]-2-ethylindolizin-1-yl]glyoxylamide 622836-45-7P, 2-[3-[(Biphenyl-2-yl)methyl]-2-ethyl-8-[[[[(1-methylcyclopentyl)oxy]carbonyl]ethyl]oxy]carbonyl]methyl]oxy]indolizin-1-yl]glyoxylamide 622836-46-8P, 2-[3-[(Biphenyl-2-yl)methyl]-2-ethyl-8-[[[[(2-morpholino)ethyl]oxy]carbonyl]methyl]oxy]indolizin-1-yl]glyoxylamide 622836-47-9P, 622836-48-0P, 2-[3-[(Biphenyl-2-yl)methyl]-2-ethyl-8-[[[[(2-oxopropyl)oxy]carbonyl]methoxy]indolizin-1-yl]glyoxylamide 622836-49-1P, 2-[3-[(Biphenyl-2-yl)methyl]-2-ethyl-8-[[[[(1-trityltetrazol-5-yl)methyl]oxy]indolizin-1-yl]glyoxylamide 622836-50-4P, 2-[7-(2-Carboethoxyethyloxy)-2-ethyl-3-(o-phenylbenzyl)indolizin-1-yl]glyoxylamide 622836-58-2P, 2-[3-(Biphenyl-2-yl)-8-[(carboxymethyl)oxy]-2-methoxyindolizin-1-yl]glyoxylamide

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(sPLA2 inhibitor; prepn. of indolizines as inhibitors of sPLA2 mediated release of fatty acids)

L5 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN

AN 1988:5780 CAPLUS

DN 108:5780

TI 6-(1-carbamoyl-1-hydroxymethyl)penicillanic acid derivatives, their preparation, and their use as antibacterial agents and/or .beta.-lactamase inhibitors

IN Barth, Wayne Ernest

PA Pfizer Inc., USA

SO Eur. Pat. Appl., 138 pp.

CODEN: EPXXDW

DT Patent




LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 220939	A1	19870506	EP 1986-308235	19861023
	R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
	WO 9006928	A1	19900628	WO 1985-US2134	19851029
	W: US				
	DK 8605143	A	19870703	DK 1986-5143	19861028
	JP 62142183	A2	19870625	JP 1986-258106	19861029
	JP 06092417	B4	19941116		
	US 4797394	A	19890110	US 1987-85675	19870605

10/620209

US 4868296 A 19890919 US 1988-243568 19880912
 PRAI WO 1985-US2134 19851029
 US 1987-85675 19870605
 OS CASREACT 108:5780
 GI




- AB Title compds. I [n = 0-2; R = H, ester group hydrolyzable under physiol. conditions, acyloxymethyl or 1-(acyloxy)ethyl derived from conventional .beta.-lactam antibiotics; R₁, R₂ = H, (un)substituted Ph, phenylalkyl, cycloalkyl, naphthyl, azolyl, etc.; NR₁R₂ = pyrrolidino, piperidino, morpholino, 1,2,3,4-tetrahydroquinolinyl, etc.] and their salts, useful as antibacterial agents and/or .beta.-lactamase inhibitors (no data), were prepd. by a) hydrogenolysis of I (R = CH₂Ph) and optionally b) converting the compd. to a cationic salt or c) converting the compd. to an acid addn. salt if the compd. contains a basic N atom. Further, the compds. may be converted to physiol. hydrolyzable esters or to acyloxymethyl or 1-(acyloxy)ethyl esters derived from conventional .beta.-lactam antibiotics. The benzyl ester was prepd. by a) reacting a cyclic anhydride II (q = 0, 2) with HNR₁R₂ and b) if desired, oxidizing the resulting 6-carbamoyl benzyl ester I (R = CH₂Ph, n = 0) to a benzyl ester (n = 1 or 2) with 1 or 2 mol equiv 3-ClC₆H₄C(O)OOH. II are prepd. by a) reacting 6-dibromo compds. III with 1 mol equiv methylmagnesium Grignard reagent and then with H₂C:CHCH₂OCOCOH to form allyl ester IV; b) debromination to give V; c) hydrolysis to give the acid VI; and d) reaction with COCl₂ in the presence of tertiary amine. Benzyl 6,6-dibromopenicillanate (III, q = 0) was treated with MeMgBr at -78.degree., then allyl glyoxalate at -78.degree. to give (R)- and (S)-IV (q = 0) the (R)-isomer of which was debrominated to give (S)-V (q = 0). Treating this with BuCH₂CO₂Na, then Pd(PPh₃)₄ gave the Na salt of (S)-VI which was successively treated with COCl₂ and NH₄OH to give (S)-I (R = CH₂Ph, R₁ = R₂ = H, n = 0). Hydrogenolysis in the presence of NaHCO₃ and 10% Pd/C gave (S)-I (R = Na, R₁ = R₂ = H, n = 0).
- IT 4706-32-5P, N-Glyoxyloypiperidine 16423-59-9P, N-Glyoxyloymorpholine 79036-50-3P, N,N-Dimethylglyoxylamide 106435-93-2P, N-Glyoxyloypyrrolidine 111605-39-1P, N-Isopropylglyoxylamide
 RL: SPN (Synthetic preparation); PREP (Preparation) (prepn. of)
- IT 75-16-1, Methylmagnesium bromide
 RL: RCT (Reactant); RACT (Reactant or reagent) (reaction of, with benzyl dibromopenicillanate and dimethylglyoxylamide)
- IT 4706-32-5 16423-59-9, N-Glyoxyloymorpholine 64370-42-9, Allyl glyoxalate 79036-50-3, N,N-Dimethylglyoxylamide 106435-93-2, N-Glyoxyloypyrrolidine 111605-39-1
 RL: RCT (Reactant); RACT (Reactant or reagent) (reaction of, with benzyl dibromopenicillanate and methylmagnesium bromide)
- IT 35564-99-9, Benzyl 6,6-dibromopenicillanate
 RL: RCT (Reactant); RACT (Reactant or reagent) (reaction of, with methylmagnesium bromide and dimethylglyoxylamide)

L5 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN
 AN 1986:552787 CAPLUS

10/620209

DN 105:152787
TI Synthesis of psilocin labeled with carbon-14 and tritium
AU Poon, Grace; Chui, Yun Cheung; Law, Francis C. P.
CS Dep. Biol. Sci., Simon Fraser Univ., Burnaby, BC, V5A 1S6, Can.
SO Journal of Labelled Compounds and Radiopharmaceuticals (1986), 23(2),
167-74
CODEN: JLCRD4; ISSN: 0362-4803
DT Journal
LA English
OS CASREACT 105:152787
GI



- AB 14C- and 3H-labeled psilocin (I, X = CH₂14CH₂; C₃H₂C₃H₂) tryptamine), the principal active agent of hallucinogenic mushrooms, was synthesized from 2-methyl-3-nitrophenol via 4-benzyloxyindole. 4-Benzyloxygramine was treated with K¹⁴CN to give 14C-4-benzyloxy-3-indoleacetic acid, an intermediate for I (X = CH₂14CH₂). LiAlH₄ was used to reduce 4-benzyloxy-3-indole-N,N-dimethylglyoxylamide to give I (X = C₃H₂C₃H₂).
- AB 14C- and 3H-labeled psilocin (I, X = CH₂14CH₂; C₃H₂C₃H₂) tryptamine), the principal active agent of hallucinogenic mushrooms, was synthesized from 2-methyl-3-nitrophenol via 4-benzyloxyindole. 4-Benzyloxygramine was treated with K¹⁴CN to give 14C-4-benzyloxy-3-indoleacetic acid, an intermediate for I (X = CH₂14CH₂). LiAlH₄ was used to reduce 4-benzyloxy-3-indole-N,N-dimethylglyoxylamide to give I (X = C₃H₂C₃H₂).

L5 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN
AN 1959:28666 CAPLUS
DN 53:28666
OREF 53:5137e-g
TI Glyoxylamide derivatives
IN Whitfield, Gordon H.
PA Imperial Chemical Industries Ltd.
DT Patent
LA Unavailable
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	GB 797604		19580702	GB	
AB	N,N-Dialkyl substituted glyoxylamide derivs., useful as herbicides were prepd. Me ₂ NCOCH(OH)NMe ₂ (I) (b ₈ 70-3.degree.) (41.2 g.) was dissolved in 50 ml. MeOH and poured into a column (1 1/2" .times. 3') packed with 500 g. polystyrenesulfonic acid cation-exchange resin, the resin washed with two 500 ml. portions MeOH, and 2 eluate fractions were collected. Removal of MeOH from the 1st eluate and distn. of the residue gave 20.69 g. N,N-dimethylglyoxylamide Me hemiacetal (II), b ₂₀ 82.86.degree.. Similar treatment of the 2nd MeOH eluate gave 3.78 g. II. Exposure of II to moist air or treatment with the theoretical amt. of H ₂ O gave Me ₂ NCOCHO.0.5H ₂ O (III), m. 121.degree.. Similar treatment of I in H ₂ O gave III, m. 121-2.degree., directly. Cf. C.A. 53, 227d.				
AB	N,N-Dialkyl substituted glyoxylamide derivs., useful as herbicides were prepd. Me ₂ NCOCH(OH)NMe ₂ (I) (b ₈ 70-3.degree.) (41.2 g.) was dissolved in 50 ml. MeOH and poured into a column (1 1/2" .times. 3') packed with 500 g. polystyrenesulfonic acid cation-exchange resin, the resin washed with two 500 ml. portions MeOH, and 2 eluate fractions were collected. Removal of MeOH from the 1st eluate and distn. of the residue gave 20.69 g. N,N-dimethylglyoxylamide Me hemiacetal (II), b ₂₀ 82.86.degree.. Similar treatment of the 2nd MeOH eluate gave 3.78 g. II. Exposure of II to moist air or treatment with the theoretical amt. of H ₂ O gave Me ₂ NCOCHO.0.5H ₂ O (III), m. 121.degree.. Similar treatment of I in H ₂ O gave III, m. 121-2.degree., directly. Cf. C.A. 53, 227d.				

L5 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN
AN 1959:1738 CAPLUS
DN 53:1738
OREF 53:227d-f
TI Glyoxylic acid derivatives

10/620209

=> s e4-e6

1 61960-32-5/BI
(61960-32-5/RN)

1 79036-50-3/BI
(79036-50-3/RN)

1 939-71-9/BI
(939-71-9/RN)


L7 3 (61960-32-5/BI OR 79036-50-3/BI OR 939-71-9/BI)

=> d scan

L7 3 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN

IN 1,3-Benzodioxole-2-carboxamide (6CI, 7CI, 8CI)

MF C8 H7 N O3




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):2

L7 3 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN

IN Acetamide, 2-hydroxy-2-methoxy-N,N-dimethyl- (9CI)

MF C5 H11 N O3




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L7 3 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN

IN Acetamide, N,N-dimethyl-2-oxo- (9CI)

MF C4 H7 N O2



10/620209

IN Whitfield, Gordon H.
PA Imperial Chemical Industries Ltd.
DT Patent
LA Unavailable


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	GB 793807		19580423	GB	
AB	<p>R1R2NCOCHO, where R1 and R2 are alkyl groups, is prepd. by treating R1R2NCO with LiC.tplbond.CH to yield R1R2NCOCH(OLi)NR1R2 followed by hydrolyzing to R1R2NCOCH(OH)NR1R2 (I). I with an acid gives R1R2NCOCHO and the acid salt of R1R2NH. E.g., 219 g. Me2NCO added dropwise to 16 g. LiC.tplbond.CH in 150 ml. boiling MePh, refluxed 0.5 hr., and MePh and unreacted Me2NCO distd. in vacuo left 108 g. Me2NCOCH(OLi)NMe2 (II). II added to 250 ml. H2O, extd. with ether, dried, and distd. yielded Me2NCOCH(OH)NMe2 (III), b8.0 70-3.degree.. A small portion of III with 2,4-dinitrophenylhydrazine sulfate yielded N,N-dimethylglyoxylamide 2,4-dinitrophenylhydrazone, m. 208.degree.. III is useful as an intermediate in the prepn. of herbicides and pharmaceuticals.</p>				
AB	<p>R1R2NCOCHO, where R1 and R2 are alkyl groups, is prepd. by treating R1R2NCO with LiC.tplbond.CH to yield R1R2NCOCH(OLi)NR1R2 followed by hydrolyzing to R1R2NCOCH(OH)NR1R2 (I). I with an acid gives R1R2NCOCHO and the acid salt of R1R2NH. E.g., 219 g. Me2NCO added dropwise to 16 g. LiC.tplbond.CH in 150 ml. boiling MePh, refluxed 0.5 hr., and MePh and unreacted Me2NCO distd. in vacuo left 108 g. Me2NCOCH(OLi)NMe2 (II). II added to 250 ml. H2O, extd. with ether, dried, and distd. yielded Me2NCOCH(OH)NMe2 (III), b8.0 70-3.degree.. A small portion of III with 2,4-dinitrophenylhydrazine sulfate yielded N,N-dimethylglyoxylamide 2,4-dinitrophenylhydrazone, m. 208.degree.. III is useful as an intermediate in the prepn. of herbicides and pharmaceuticals.</p>				

10/620209

=> d scan


L6 3 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN
IN Glycolamide, 2-dimethylamino-N,N-dimethyl- (6CI)
MF C6 H14 N2 O2
CI COM



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):2

L6 3 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN
IN Acetic acid, oxo- (9CI)
MF C2 H2 O3
CI COM



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L6 3 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN
IN Acetamide, 2-[(2,4-dinitrophenyl)hydrazono]-N,N-dimethyl- (9CI)
MF C10 H11 N5 O5



10/620209

=> d his

(FILE 'HOME' ENTERED AT 17:57:21 ON 16 JAN 2004)

FILE 'CAPLUS' ENTERED AT 18:00:09 ON 16 JAN 2004

E SCHLOEMER G/IN


L1	17 S E4-E6
L2	0 S L1 AND ACETAMIDE?
L3	0 S L1 AND IMIDAZO?
L4	7 S L1 AND PROCESS

=>

10/620209

L4 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2004 ACS on STN
 AN 2003:222366 CAPLUS
 DN 138:238439
 TI Production of benazepril and analogs by kinetic resolution of an intermediate
 IN Tseng, Wei-Hong; Cheng, Kau-Ming; Schloemer, George; Chen, Chien-Wen; Cheng, Chih-Wen
 PA Scinopharm Taiwan, Ltd., Taiwan
 SO U.S. Pat. Appl. Publ., 7 pp., Cont.-in-part of U.S. Ser. No. 910,509.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2003055245	A1	20030320	US 2002-151772	20020521
	US 2002183515	A1	20021205	US 2001-910509	20010719
	US 6548665	B2	20030415		
PRAI	US 2001-291888P	P	20010518		
	US 2001-910509	A2	20010719		
OS	MARPAT 138:238439				
GI					



AB A process for the prepn. of benazepril and analogs I by reaction of compd. II [Z₁ = halogen] with compd. III [R₁ = H, alkyl or a combination of H and alkyl; R₂ = alkyl] in a polar solvent via epimerization and kinetic resolu. of intermediate catalyzed by phase transfer catalyst was developed. Thus, coupling of L-homophenylalanine Et ester to 3-bromo-2,3,4,5-tetrahydro-1H-1-benzapin-2-one using sodium iodide, epimerization and kinetic resolu. of intermediate carboxylic acid, followed by esterification gave compd. (S,S)-I (R₁ = H, R₂ = Et) in 80% yield and the ratio of enantiomers detd. by HPLC is SS:RR > 99.5:0.5.

=> d 2-7 bib abs

L4 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2004 ACS on STN
 AN 2002:736940 CAPLUS
 DN 137:263201
 TI Process for making taxane derivatives
 IN Schloemer, George; Chen, Yung-fa; Lin, Chien Hsin; Daniewski, Wlodzimierz
 PA Scinopharm Taiwan, Ltd., Taiwan
 SO U.S. Pat. Appl. Publ., 9 pp.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002137955	A1	20020926	US 2001-815517	20010323
	US 6531611	B2	20030311		
	WO 2002076967	A1	20021003	WO 2001-US9348	20010323
	W:	CN, JP			
	RW:	AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR			

10/620209

PRAI US 2001-815517 A 20010323
OS CASREACT 137:263201; MARPAT 137:263201
GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The present invention provides a novel semi-synthetic method of producing a variety of novel taxane derivs. I (R1 = alkoxy, R2 = alkoxy, H; R3 = alkyl; R4 = alkyl, aryl; X = protective group) by reaction of a phenylisoserine deriv. II with a suitably blocked Baccatin III deriv. III. I may be further modified to form paclitaxel and other potentially useful taxane derivs. Thus, III (R1 = R2 = MeO; R3 = Me; R4 = Ph), prepd. from (2R,3S)-phenylisoserine-HCl and .alpha.-methylcinnamic acid, was treated with 7-triethylsilylbaccatin III to give the corresponding I, which was converted to paclitaxel in 4 steps.

L4 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2004 ACS on STN
AN 2002:290826 CAPLUS
DN 136:310051
TI Process for the preparation of 4,4'-diketo-.beta.-carotene derivatives
IN Schloemer, George C.; Schloemer, Danuta A.; Davis, Jeffery L.
PA Prodemex, S.A. D.E.C.V., Mex.
SO U.S., 4 pp.
CODEN: USXXAM
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6372946	B1	20020416	US 2001-953007	20010913
	NO 2002004266	A	20030314	NO 2002-4266	20020906
	CN 1417207	A	20030514	CN 2002-141620	20020906
	EP 1293499	A1	20030319	EP 2002-256236	20020909
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				

PRAI US 2001-953007 A 20010913

OS CASREACT 136:310051

AB A method of prepg. .beta.-carotene derivs. such as canthaxanthin and astaxanthin was described. The method employs an in situ system to generate hypobromous acid as the oxidizing agent using a salt of sulfite, hydrogen sulfite or bisulfite in combination with a bromate salt. Astaxanthin and canthaxanthin were obtained in good yield with a significantly reduced reaction time. Thus, zeaxanthin was oxidized using sodium hydrogen sulfite in chloroform to form axtaxanthin.

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2004 ACS on STN
AN 2001:798186 CAPLUS
DN 135:344616
TI Oxidative process for the preparation of astaxanthin from zeaxanthin using a halogenating agent with the salt of chloric or bromic acid in an inert solvent
IN Schloemer, George C.; Davis, Jeffery L.
PA Prodemex, S.A. de C.V., Mex.
SO PCT Int. Appl., 12 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001081301	A2	20011101	WO 2001-US13295	20010425
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, EE, ES, FI, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	US 2001051357	A1	20011213	US 2001-813685	20010319

10/620209

US 6376717 B2 20020423
EP 1276719 A2 20030122 EP 2001-932633 20010425
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR
NO 2001006293 A 20020211 NO 2001-6293 20011220
ZA 2001010503 A 20020829 ZA 2001-10503 20011221
PRAI US 2000-199875P P 20000426
US 2001-813685 A 20010319
WO 2001-US13295 W 20010425
OS CASREACT 135:344616
AB Astaxanthin is prepd. from zeaxanthin by oxidn. using a halogenating agent with the salt of chloric or bromic acid in an inert solvent.

L4 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2004 ACS on STN
AN 1996:689881 CAPLUS
DN 126:19174
TI Preparation of acyclovir using 1,3 dioxolane
IN Schloemer, George C.; Han, Yeun-kwei; Harrington, Peter J.
PA Syntex (U.S.A.) Inc., USA
SO U.S., 8 pp., Cont.-in-part of U.S. Ser. No. 280,269, abandoned.
CODEN: USXXAM
DT Patent
LA English
FAN.CNT 2


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 5567816	A	19961022	US 1995-426005	19950427
	CA 2152863	AA	19960127	CA 1995-2152863	19950628
	JP 08053451	A2	19960227	JP 1995-176022	19950712
	EP 709385	A1	19960501	EP 1995-110955	19950713
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
	CN 1122805	A	19960522	CN 1995-115316	19950725
	BR 9503442	A	19960604	BR 1995-3442	19950725
	FI 9503580	A	19960127	FI 1995-3580	19950726
PRAI	US 1994-280269		19940726		
	US 1995-426005		19950427		

AB A process for the prepn. of acyclovir via coupling of guanine or silylated guanines with 1,3-dioxolane in the presence of a selective alkylation catalyst selected from the group consisting of trifluoromethanesulfonic acid, trimethylsilyl trifluoromethanesulfonate, and bistrimethylsilyl sulfonate, and hydrolyzing the product thus formed.

L4 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2004 ACS on STN
AN 1996:313501 CAPLUS
DN 124:343989
TI Method for producing 9-(2-hydroxyethoxymethyl)guanine (acyclovir) as antiviral agent
IN Han, Yuen-Kwei; Harrington, Peter John; Schloemer, George Charles
PA F. Hoffmann-La Roche Ag, Switz.
SO Can. Pat. Appl., 28 pp.
CODEN: CPXXEB
DT Patent
LA English
FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CA 2152863	AA	19960127	CA 1995-2152863	19950628
	US 5567816	A	19961022	US 1995-426005	19950427
PRAI	US 1994-280269		19940726		
	US 1995-426005		19950427		

OS CASREACT 124:343989; MARPAT 124:343989
GI



AB An efficient and selective process for the synthesis of the antiviral 9-(2-hydroxyethoxymethyl)guanine (acyclovir) (I) involves (1) contacting a silylated guanine or mixts. of silylated guanine (II; Z1, Z2, Z3 = H, R1R2R3Si; wherein R1 - R3 = lower alkyl; provided that at least one of Z1 - Z3 = R1R2R3Si) with 1,3-dioxolane (III) in the presence of a selective alkylation catalyst and (2) hydrolyzing the product formed. Said catalyst is selected from CF3SO3H, CF3SO3SiMe3, and bis(trimethylsilyl) sulfonate and CF3SO3SiMe3 is generated by contacting CF3SO3H with hexamethyldisilazane. The process avoids the use of acyl groups for protection of guanine, essentially specific for the prepn. of the N-9 isomer, thus eliminates the need for the chromatog. sepn. of the N-9/N-7 isomer mixts., provides I in good yields, requires simple starting materials and reaction conditions, and is carried out from start to finish in a single reaction vessel. Thus, a mixt. of 25 g guanine, 125 mL hexamethyldisilazane, and 1 mL CF3SO3SiMe3 was refluxed at 130-135.degree. for 24 h, cooled to 70.degree., treated with 25 mL 1,3-dioxolane, refluxed for 16 h, distd. under reduced pressure to remove excess hexamethyldisilazane, cooled to 70.degree., poured into a mixt. of 600 mL 10% aq. AcOH, and heated to give a soln. The soln. was treated with 1.25 g activated carbon to remove any color, filtered, and the filtrate was slowly cooled to 5.degree. to give, after filtering off the white cryst. solid formed, 78% I.

L4 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2004 ACS on STN

AN 1989:23725 CAPLUS

DN 110:23725

TI Process for preparing (+-)-1,2-dihydro-3H-pyrrolo[1,2-
a]pyrrole-1,7-dicarboxylates as intermediates for pharmaceuticals

IN Khatri, Hiralal N.; Fleming, Michael P.; Schloemer, George C.

PA Syntex (U.S.A.), Inc., USA

SO Eur. Pat. Appl., 28 pp.

CODEN: EPXXDW

DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 275092	A1	19880720	EP 1988-100390	19880113
	EP 275092	B1	19920603		
	R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
	US 4835288	A	19890530	US 1987-3162	19870114
	US 4849526	A	19890718	US 1987-3104	19870114
	DK 8800143	A	19880715	DK 1988-143	19880113
	FI 8800133	A	19880715	FI 1988-133	19880113
	FI 90344	B	19931015		
	FI 90344	C	19940125		
	NO 8800127	A	19880715	NO 1988-127	19880113
	NO 169124	B	19920203		
	NO 169124	C	19920513		
	AU 8810240	A1	19880721	AU 1988-10240	19880113
	AU 613334	B2	19910801		
	JP 63198684	A2	19880817	JP 1988-6757	19880113
	ZA 8800225	A	19890927	ZA 1988-225	19880113
	HU 48881	A2	19891128	HU 1988-117	19880113
	HU 200606	B	19900728		
	HU 51595	A2	19900528	HU 1989-5354	19880113
	HU 201728	B	19901228		
	HU 52045	A2	19900628	HU 1989-5355	19880113
	HU 203532	B	19910828		
	HU 52046	A2	19900628	HU 1989-5356	19880113

10/620209

HU 203721	B	19910930		
IL 85094	A1	19910916	IL 1988-85094	19880113
IL 96388	A1	19910916	IL 1988-96388	19880113
IL 96389	A1	19910916	IL 1988-96389	19880113
AT 76873	E	19920615	AT 1988-100390	19880113
ES 2041703	T3	19931201	ES 1988-100390	19880113
HU 213614	B	19970828	HU 1990-1266	19880113
CA 1340404	A1	19990223	CA 1988-556465	19880113
US 4874872	A	19891017	US 1988-255799	19881011
US 4937368	A	19900626	US 1989-299701	19890123
NO 9003993	A	19880715	NO 1990-3993	19900913
NO 174583	B	19940221		
NO 174583	C	19940601		
NO 9003994	A	19880715	NO 1990-3994	19900913
NO 174346	B	19940110		
NO 174346	C	19940420		
NO 9003995	A	19880715	NO 1990-3995	19900913
NO 173828	B	19931101		
NO 173828	C	19940209		
FI 92488	B	19940815	FI 1991-2709	19910605
FI 92488	C	19941125		
FI 95242	B	19950929	FI 1991-2710	19910605
FI 95242	C	19960110		
FI 91148	B	19940215	FI 1993-320	19930126
FI 91148	C	19940525		
PRAI US 1987-3104	A	19870114		
US 1987-3162	A	19870114		
EP 1988-100390	A	19880113		
HU 1988-117	A	19880113		
IL 1988-85094	A	19880113		
NO 1988-127	A1	19880113		
OS				
GI				
			CASREACT 110:23725; MARPAT 110:23725	



AB A process for producing diesters I (R = alkyl), useful as intermediates for pharmaceuticals II [Ar = alkyl, alkoxy, or halo (un)substituted Ph, 2- or 3- furoyl, 2- or 3-thienyl, 2- or 3-pyrrolyl; R = H, alkyl] useful as analgesics, antiinflammatories, antipyretics, and smooth muscle relaxants (no data), comprised: a) cyclizing pyrrole III (X = halo) with a hindered amine in an aprotic polar solvent; or b) reacting pyrrolidine IV with XCH2CHO (X = halo) in aq. soln. I' (R = alkyl) are sapond. to I (R = H) which are monoesterified to I (R at 1 = alkyl, R at 7 = H) which are decarboxylated to II (no ArCO group). These are aroylated with an amide or morpholide to give II. I (R = H), which had been prepd. in 5 steps from BrCH2CH2NH2.HBr and (MeO2CCH2)2CO was converted in 4 steps into II (Ar = 4-MeC6H4, R = H).

L Number	Hits	Search Text	DB	Time stamp
1	3478	phosphorus adj tribromide	USPAT; US-PGPUB	2004/01/16 17:19
2	30687	thionyl adj chloride	USPAT; US-PGPUB	2004/01/16 17:20
3	315	((phosphorus adj tribromide) near (thionyl adj chloride))	USPAT; US-PGPUB	2004/01/16 17:20
4	593	imidazopyridine	USPAT; US-PGPUB	2004/01/16 17:21
5	0	((phosphorus adj tribromide) near (thionyl adj chloride)) and imidazopyridine	USPAT; US-PGPUB	2004/01/16 17:20
6	271	imidazopyridines	USPAT; US-PGPUB	2004/01/16 17:21
7	745	imidazopyridine or imidazopyridines	USPAT; US-PGPUB	2004/01/16 17:21
8	0	(imidazopyridine or imidazopyridines) and ((phosphorus adj tribromide) near (thionyl adj chloride))	USPAT; US-PGPUB	2004/01/16 17:22
9	11485	halogenating	USPAT; US-PGPUB	2004/01/16 17:22
10	0	((phosphorus adj tribromide) near (thionyl adj chloride)) near halogenating	USPAT; US-PGPUB	2004/01/16 17:22
11	180	((phosphorus adj tribromide) near (thionyl adj chloride)) same halogenating	USPAT; US-PGPUB	2004/01/16 17:45
12	3	((((phosphorus adj tribromide) near (thionyl adj chloride)) same halogenating) and sleep	USPAT; US-PGPUB	2004/01/16 17:47
13	113	hydrolysis and (((phosphorus adj tribromide) near (thionyl adj chloride)) same halogenating)	USPAT; US-PGPUB	2004/01/16 17:47
14	310	hydrolysis same (phosphorus adj tribromide) (((phosphorus adj tribromide) near (thionyl adj chloride)) same halogenating)	USPAT; US-PGPUB	2004/01/16 17:47
15	1	hydrolysis same (((phosphorus adj tribromide) near (thionyl adj chloride)) same halogenating)	USPAT; US-PGPUB	2004/01/16 17:48